

Infant Mortality and Its Predictors in Benshangul-Gumuz Region, Ethiopia: Further Analysis of 2019 Ethiopian Mini Demographic and Health Survey

Temesgen Senbeto^{1,*}, Gizachew Gobebo¹, Woldemariam Erkalo²

¹College of Natural and Computational Science, Ambo University, Ambo, Ethiopia

²College of Natural and Computational Science, Mettu University, Mettu, Ethiopia

Email address:

temesenbeto@gmail.com (Temesgen Senbeto)

*Corresponding author

To cite this article:

Temesgen Senbeto, Gizachew Gobebo, Woldemariam Erkalo. Infant Mortality and Its Predictors in Benshangul-Gumuz Region, Ethiopia: Further Analysis of 2019 Ethiopian Mini Demographic and Health Survey. *Journal of Health and Environmental Research*. Vol. 9, No. 3, 2023, pp. 90-94. doi: 10.11648/j.jher.20230903.13

Received: August 21, 2023; Accepted: September 18, 2023; Published: October 14, 2023

Abstract: Background: Infant mortality rate has dramatically been fallen in the last two decades globally. Ethiopia, with infant mortality rate of 34 deaths per 1000 live births, is among the countries with high infant mortality in Africa. The main purpose of this study was to investigate predictors of infant mortality in Benshangul-Gumuz region, Ethiopia. Methods: The source of data for this study was 2019 Ethiopian Mini Demographic and Health Survey (EMDHS). Multivariable logistic regression model was used to identify the predictors of infant mortality. Results: Multiple births and infants born from mothers aged 34 years and older were at higher risk of dying before celebrating their first birthdays. whereas females, infants born from mothers with secondary and higher education, infants born from mothers who had ANC visits of at least one time, and infants born at health institution were at lower risk of dying before celebrating their first birthdays. Conclusions: Infant mortality rate in region was high. Birth type, sex of child, maternal age, maternal educational level, place of delivery, and antenatal care visits were predictors of infant mortality in Benshangul-Gumuz region, Ethiopia. Hence, health institutional delivery is recommended in the region to minimize the infant mortality rate in the region.

Keywords: Predictors, Infant Mortality, Benshangul-Gumuz Region, Ethiopia

1. Introduction

Infant mortality rate has dramatically been fallen in the last two decades globally [1, 2]. Despite the dramatic decline in the infant mortality rate globally, low-and middle-income countries still share the largest burden of infant mortality [3-5]. Ethiopia, with infant mortality rate of 34 deaths per 1000 live births in 2021 [6], is among the countries with high infant mortality in Africa.

In spite of the reduction in infant mortality rate in Ethiopia over the past 14 years from 74 in 2005 to 47 in 2019 [7], it increased from 62 in 2016 [8] to 74 in 2019 [7] in Benshangul-Gumuz region in Ethiopia.

Region, residence, religion, employment status of mother, economic status, father's educational level, educational level, maternal age, mother's age at first birth, mother's marital

status, maternal anaemia, HIV status of mother, pregnancy type, diarrhoea, source of drinking water, birth type, parity, sex of child, birth order, place of delivery, body mass index, ANC visit and preceding birth interval were factors associated with infant mortality [9-36].

Benshangul-Gumuz region of Ethiopia with the infant mortality rate of 74, which is very high figure, is the region with the highest infant mortality in the country in 2019 [7]. Thus, the main objective of this study was to investigate predictors of infant mortality in Benshangul-Gumuz region, Ethiopia.

2. Methods

2.1. Data

The source of data for this study was 2019 EMDHS data.

A total of 530 live births were included in the study.

2.2. Variables of the Study

2.2.1. Response Variable

Response variable is the death of child before celebrating first birthday.

2.2.2. Explanatory Variables

The Explanatory variables were place of residence, sex of child, birth order, type of birth, parity, maternal educational level, maternal age, marital status of mother, household wealth index, ANC visits, maternal age at first birth, place of delivery, mode of delivery and source of drinking water.

2.3. Data Analysis

Multivariable logistic regression analysis was done to investigate the risk factors of infant mortality. The data were analyzed by SPSS version 26.

3. Results

3.1. Socioeconomic Characteristics and Results of Bivariate Analysis

Among the total live-born children included, majority of the infants 461 (87.0%) were born in rural areas. About 52.1 % were males and 47.9% were females. About half

(49.4%) had birth order of 4th and above. With regard to type of birth, majority (94.3%) had birth type of single and only 5.7% had multiple birth type. About 41.7%, 30.6% and 27.7% were born from families with total children ever born of 5 and more, 1 to 2, and 3 to 4 respectively. More than half (52.1%) were born from uneducated mothers. Majority (72.5%) were born from mothers aged less than 25 years. More than two-fifth (47.5%) were born from mothers whose age at first birth was less than 20 years. About 61% of the children were born from mothers from poor families. More than three- fourth (86.6%) of mothers attended the ANC follow-up during their pregnancies at least one time. About two-third (66.0%) of the children were born at health institutions. Majority (95.8%) of the children were born vaginally. About four-fifth (79.4%) were born from families who use protected source of drinking water (Table 1).

Of the total children, 43 (8.1%) died before their first birth day. The percentage of infant mortality was higher (8.7%) among rural children than urban ones (4.3%). The highest percentage of infant mortality (10.6%) was among children whose mothers had primary education. The higher percentage (10.6%) of the infant mortality was also observed among children born at home than children born at health institutions (6.9%). The higher percentage (8.5%) of the infant mortality was observed among children whose mothers did not attend the ANC visit (Table 1).

Table 1. Socioeconomic characteristics of the participants and bivariate analysis result of infant mortality in Benshangul-Gumuz region.

Variables	Count	Percent	Child died before celebrating first birthday		P-value
			Yes (%)	No (%)	
Residence					
Urban	69	13.0	3 (4.3)	66 (95.7)	0.044*
Rural	461	87.0	40 (8.7)	421 (91.3)	
Sex of child					
Male	276	52.1	24 (8.7)	252 (91.3)	0.002*
Female	254	47.9	19 (7.5)	235 (92.5)	
Birth order					
First	103	19.4	12 (11.7)	91 (88.3)	0.190
Second -Third	165	31.1	9 (5.5)	156 (94.5)	
Fourth and above	262	49.4	22 (8.4)	240 (91.6)	
Birth type					
Single	500	94.3	30 (6.0)	470 (94.0)	0.000*
Multiple	30	5.7	13 (43.3)	17 (56.7)	
Parity					
1-2	162	30.6	14 (8.6)	148 (91.4)	0.049*
3-4	147	27.7	7 (4.8)	140 (95.2)	
5 and more	221	41.7	22 (10.0)	199 (90.0)	
Maternal education					
No education	276	52.1	19 (6.9)	257 (93.1)	0.032*
Primary	208	39.2	22 (10.6)	186 (89.4)	
Secondary and higher	46	8.7	2 (4.3)	44 (95.7)	
Maternal age					
Less than 25	384	72.5	28 (7.3)	356 (92.3)	0.029*
25-34	35	6.6	7 (20.0)	28 (80.0)	
34 and above	111	20.9	8 (7.2)	103 (92.8)	
Mother's age at first birth					
Less than 20	252	47.5	14 (5.6)	238 (94.4)	0.041*
20-24	132	24.9	14 (10.6)	118 (89.4)	
25 and above	146	27.5	15 (10.3)	131 (89.7)	
Marital status of mother					
Never in union	2	0.4	0 (0.0)	2 (100.0)	0.555

Variables	Count	Percent	Child died before celebrating first birthday		P-value
			Yes (%)	No (%)	
Currently in union	497	93.8	39 (7.8)	458 (92.2)	0.998
Widowed/divorced/separated	31	5.8	4 (12.9)	27 (87.1)	
Household wealth index					
Poor	321	60.6	26 (8.1)	295 (91.9)	0.998
Medium	109	20.6	9 (8.3)	100 (91.7)	
Rich	100	18.9	8 (8.0)	92 (92.0)	
Source of drinking water					0.130
Protected	421	79.4	38 (9.0)	383 (91.0)	
Unprotected	109	20.6	5 (4.6)	104 (95.4)	
ANC					0.012*
No	71	13.4	6 (8.5)	65 (91.5)	
Yes	459	86.6	37 (8.1)	422 (91.9)	
Mode of delivery					0.531
Vaginal	508	95.8	42 (8.3)	466 (91.7)	
Caesarean	22	4.2	1 (4.5)	21 (95.5)	
Place of delivery					0.040*
Home	180	34.0	19 (10.6)	161 (89.4)	
Health institution	350	66.0	24 (6.9)	326 (93.1)	

*: statistically significant

3.2. Predictors of Infant Mortality in Benshangul-Gumuz Region of Ethiopia

This study revealed that sex of child, type of birth, education status of mother, age of mother, ANC visits and place of delivery were risk factors associated with infant mortality in Benshangul-Gumuz region of Ethiopia. The odds of infant mortality among females was 0.690 (AOR: 0.690, 95% CI: 0.342, 0.899) times lower than males. The odds of infant mortality among multiple births was 3.067 (AOR: 3.067, 95% CI: 2.313, 10.139) times higher than among singletons. The odds of infant mortality for those born from mothers with secondary and higher education was 0.460 (AOR: 0.460, 95% CI: 0.287, 0.885) times lower than among infants born from mother with no education. The odds of infant mortality for those born from mothers who had ANC visits was 0.597 (AOR: 0.597, 95% CI: 0.326, 0.709) times lower than among infants born from mothers who had not ANC visits. Similarly, the odds of infant mortality for those born in health institution was 0.611 (AOR: 0.611, 95% CI: 0.0294, 0.0899) times lower than among infants born from mothers who had not ANC visits. The odds of infant mortality for those born from mothers were older than 34 years was 1.539 (AOR: 1.539, 95% CI: 1.183, 9.802) times higher than among infants born from mother were younger than 25 years (Table 2).

Table 2. Predictors of infant mortality in Benshangul-Gumuz region, Ethiopia.

Variables	AOR	95% CI of AOR	P-value
Residence			
Urban	Ref		
Rural	2.208	0.580, 8.339	0.245
Sex of child			
Male	Ref		
Female	0.690	0.342, 0.899	0.038*
Birth type			
Single	Ref		
Multiple	3.067	2.313, 10.139	0.000*

Variables	AOR	95% CI of AOR	P-value
Parity			
1-2	Ref		
3-4	1.588	1.204, 1.799	0.327
5 and more	1.380	0.244, 5.502	0.221
Maternal age			
Less than 25	Ref		
25-34	1.009	0.837, 3.526	0.921
34+	1.539	1.183, 9.802	0.040*
Maternal education			
No education	Ref		
Primary	0.292	0.017, 0.619	0.482
Secondary and higher	0.460	0.287, 0.885	0.007*
Mother's age at first birth			
Less than 20	Ref		
20-24	1.813	0.748, 4.391	0.188
25+	1.354	0.503, 3.648	0.549
Place of delivery			
Home	Ref		
Health institution	0.611	0.294, 0.899	0.024*
ANC			
No	Ref		
Yes	0.597	0.326, 0.709	0.033*

AOR: Adjusted odds ratio, Ref: Reference group, *: statistically significant

4. Discussion

Among the total children include in the study, 43 (8.1%) died before the first birth day. The study revealed that females were less likely to die before celebrating their first birthdays than males. This result agrees with the result of the studies [37, 38]. Multiple births were at higher risk of dying before first birthday than singletons. This result agrees with the studies [29, 39, 40]. Children born from mothers who were 34 years and older were more likely to die before first birthday compared to those born from mothers who were younger than 25 years. Children born from mothers with secondary and higher education were less likely to die before first birthday than those born from mothers with no education. This result agrees with the studies [40-43]. Children born at health institution were less likely to die before celebrating their first

birthdays than infants born at home. This result agrees with the studies [10, 42]. Children born from mothers with ANC visits were less likely to die than those born from mothers with no ANC visits. This result agrees with the studies [40, 44, 45].

5. Conclusions

In this study, birth type, sex of child, maternal age, maternal educational level, place of delivery, and antenatal care visits were predictors of infant mortality in Benshangul-Gumuz region of Ethiopia. Hence, health institutional delivery is recommended in the region to minimize the infant mortality rate in the region.

Competing Interests

Authors declare that they have no competing interests.

References

- [1] World Health Organization. WHO Global Health Observatory (GHO) data global on maternal health indicators, 2017 update [internet]. Geneva: World Health Organization; 2017.
- [2] UNICEF. 's Division of Policy and Strategy. Committing to child survival: a promise renewed. Progress report. 2012.
- [3] Lopez AD. Levels & Trends in Child Mortality: Report 2014, Estimates Developed by the UN Inter-agency Group for Child Mortality Estimation. United Nations Inter-agency Group for Child Mortality Estimation (UN IGME); 2014.
- [4] Sartorius BK, Sartorius K. Global infant mortality trends and attributable determinants—an ecological study using data from 192 countries for the period 1990–2011. *Population Health Metrics*. 2014 Dec; 12 (1): 1-5.
- [5] Ester PV, Torres A, Freire JM, Hernández V, Gil Á. Factors associated to infant mortality in Sub-Saharan Africa. *Journal of Public Health in Africa*. 2011 Sep 5; 2 (2).
- [6] United Nations Inter-agency Group for Child Mortality Estimation. Levels & trends in child mortality: report 2022: Estimates developed by the UN Inter-Agency Group for Child Mortality Estimation. United Nations Children's Fund, New York, 2023.
- [7] Ethiopian Public Health Institute (EPHI) [Ethiopia] and ICF. Ethiopia Mini Demographic and Health Survey 2019: Final Report. Rockville: EPHI and ICF; 2021.
- [8] Central Statistical Agency (CSA) [Ethiopia] and ICF. Ethiopia Demographic and Health Survey 2016. Rockville: Addis Ababa; 2016.
- [9] Kiross GT, Chojenta C, Barker D, Loxton D. Individual-, household-and community-level determinants of infant mortality in Ethiopia. *PloS one*. 2021 Mar 12; 16 (3): e0248501.
- [10] Adebe KL, Wake SK, Yadata SD, Gondol KB, Mekebo GG, Wolde TS, Bechera T, Regesa BH, Galdassa A, Yadata KD. Understanding correlates of infant mortality in Ethiopia using 2019 Ethiopian mini demographic and health survey data. *Annals of Medicine and Surgery*. 2023 May; 85 (5): 1796.
- [11] Gobebo G, Mulugeta W, Yaekob T. Determinants of women unemployment: Evidence from Ethiopia (Case of HalabaTown, SNNPR). *International Journal of Development Research*. 2017; 7 (11): 16630-9.
- [12] Kefale B, Bedada D, Negash Y, Gobebo G. Determinants of Diarrhea among Children under Age Five Using Generalized Linear Model with Bayesian Approach: The Case of Kuyu General Hospital, Oromia Region, Ethiopia. *Clinics Mother Child Health*. S. 2021; 18: 11.
- [13] Tesema GA, Seretew WS, Worku MG, Angaw DA. Trends of infant mortality and its determinants in Ethiopia: mixed-effect binary logistic regression and multivariate decomposition analysis. *BMC Pregnancy and Childbirth*. 2021 Dec; 21 (1): 1-6.
- [14] Gobebo G. Determinant factors of under-five mortality in Southern Nations, Nationalities and People's region (SNNPR), Ethiopia. *Italian journal of pediatrics*. 2021 Dec; 47 (1): 1-9.
- [15] Abate MG, Angaw DA, Shaweno T. Proximate determinants of infant mortality in Ethiopia, 2016 Ethiopian demographic and health surveys: results from a survival analysis. *Archives of Public Health*. 2020 Dec; 78 (1): 1-0.
- [16] Mekebo GG, Argawu AS, Likassa HT, Ayele W, Wake SK, Bedada D, Hailu B, Senbeto T, Bedane K, Lulu K, Daraje S. Factors influencing exclusive breastfeeding practice among under-six months infants in Ethiopia. *BMC Pregnancy and Childbirth*. 2022 Dec; 22 (1): 1-0.
- [17] Mukherjee A, Bhattacharjee S, Dasgupta S. Determinants of infant mortality in rural India: An ecological study. *Indian journal of public health*. 2019 Jan 1; 63 (1): 27.
- [18] Argawu AS, Mekebo GG. Risk factors of under-five mortality in Ethiopia using count data regression models, 2021. *Annals of Medicine and Surgery*. 2022 Sep 22: 104764.
- [19] Jain AK. Determinants of regional variations in infant mortality in rural India. *Population studies*. 1985 Nov 1; 39 (3): 407-24.
- [20] Handiso A, Mekebo GG, Galdassa A. Trends and determinants of road traffic accident human death in Kembata Tembaro zone, SNNPR, Ethiopia. *Sci J Appl Math Stat*. 2022; 10: 85-9.
- [21] Dwivedi SN, Begum S, Dwivedi AK, Pandey A. Determinants of infant mortality in rural India: A three-level model. *Health*. 2013 Nov 11; 2013.
- [22] Wolde TS, Mekebo GG. Unintended pregnancy and associated factors among pregnant women in rural Ethiopia. *Journal of Pharmaceutical Research International*. 2021; 33 (60B): 2432-40.
- [23] Kananura RM, Tetui M, Mutebi A, Bua JN, Waiswa P, Kiwanuka SN, Ekirapa-Kiracho E, Makumbi F. The neonatal mortality and its determinants in rural communities of Eastern Uganda. *Reproductive health*. 2016 Dec; 13 (1): 1-9.
- [24] Wake SK, Zewotir T, Mekebo GG, Fissuh YH. Rural-urban differentials in child body mass index over time. *BMC pediatrics*. 2023 Dec; 23 (1): 1-9.
- [25] Argawu AS, Mekebo GG, Bedane K, Makarla RK, Kefale B, Senbeto T, Lemesa R, Bechera T, Lulu K, Galdassa A, Tadesse G. Prevalence and determinants of anaemia among reproductive-aged women in Ethiopia: a nationally representative cross-sectional study. *Journal of Pharmaceutical Research International*. 2021; 33 (59B): 687-98.

- [26] Mfateneza E, Rutayisire PC, Biracyaza E, Musafiri S, Mpabuka WG. Application of machine learning methods for predicting infant mortality in Rwanda: analysis of Rwanda demographic health survey 2014–15 dataset. *BMC Pregnancy and Childbirth*. 2022 Dec; 22 (1): 1-3.
- [27] Kefale BA, Woya AA, Tekile AK, Bantie GM. Spatial Distribution of Infant Mortality in Ethiopia: Using Demographic Health Survey, the Ethiopia.
- [28] Biracyaza E, Habimana S. Modeling the Risks Factors Associated with Infant Mortality in Rwanda from 2011 to 2015: Analysis of Rwanda Demographic and Health Survey (RDHS) 2014/2015. *J. Pediatrics Child Heal. Care*. 2019; 4 (1): 1026.
- [29] Regesa BH, Senbeto T, Gobebo G, Lemesa R, Makarla RK, Tadesse G, Lulu K, Bechera T, Daraje S, Galdassa A, Bedane K, Siffir A, Olani J, Teferi G, Mossisa M, Chimdi E, Degaga GT, Woldeyohannes B, Kefale B, Hemba G. Analysis of Determinants of Neonatal Mortality in Afar and Somalia Regions, Ethiopia. *Journal of Pharmaceutical Research International*. 2022; 34 (1B): 63-71.
- [30] Adebe KL, Wake SK, Mekebo GG, Dessalegn RC, DarajeYadata S, Wolde TS, Gondol KB, Galdassa A, Amega GA, Gefela TY. Analysis of regional heterogeneity and determinants of perinatal mortality in Ethiopia. *Annals of Medicine and Surgery*. 2023 Apr; 85 (4): 902.
- [31] Gobebo G, Siffir A, Hailu B, Woldeyohannes B, Kefale B, Senbeto T, Handiso A, Reddy OC. Influencing Factors of Mortality Among Adult HIV Patients Under Antiretroviral Therapy: The Case of Hossana Queen Elleni Mohammad Memorial Hospital, Ethiopia. *Science*. 2021 Aug 24; 10 (3): 72-7.
- [32] Kittur I. Factors influencing infant mortality in urban Kenya (Doctoral dissertation, University of Nairobi).
- [33] Omariba D, Beaujot R, Rajulton F. Determinants of infant and child mortality in Kenya: an analysis controlling for frailty effects. *Population Research and Policy Review*. 2007 Jun; 26 (3): 299-321.
- [34] Wake SK, Mekebo GG, Fissuh YH. Children body mass index trends in four low-and middle-income countries: Analysis of young lives cohort study data. *Clinical Epidemiology and Global Health*. 2023 Sep 1; 23: 101378.
- [35] Woldeyohannes B, Yohannes Z, Likassa HT, Mekebo GG, Wake SK, Sisay AL, Gondol KB, Argaw A, Diriba G, Irana T. Count regression models analysis of factors affecting antenatal care utilization in Ethiopia. *Annals of Medicine and Surgery*. 2023 Oct 1; 85 (10): 4882-6.
- [36] Mugo NS, Agho KE, Zwi AB, Damundu EY, Dibley MJ. Determinants of neonatal, infant and under-five mortality in a war-affected country: analysis of the 2010 Household Health Survey in South Sudan. *BMJ global health*. 2018 Feb 1; 3 (1): e000510.
- [37] Negera A, Abelti G. An analysis of the trends, differentials and key proximate determinants of infant and under-five mortality in Ethiopia: further analysis of the 2000, 2005, and 2011 demographic and health surveys.
- [38] Titaley CR, Dibley MJ, Agho K, Roberts CL, Hall J. Determinants of neonatal mortality in Indonesia. *BMC public health*. 2008 Dec; 8 (1): 1-5.
- [39] Lau C, Ambalavanan N, Chakraborty H, Wingate MS, Carlo WA. Extremely low birth weight and infant mortality rates in the United States. *Pediatrics*. 2013 May; 131 (5): 855-60.
- [40] Fenta SM, Fenta HM. Risk factors of child mortality in Ethiopia: application of multilevel two-part model. *PloS one*. 2020 Aug 17; 15 (8): e0237640.
- [41] Rahman A, Hossain Z, Kabir E, Rois R. Machine Learning Algorithm for Analysing Infant Mortality in Bangladesh. In *International Conference on Health Information Science* 2021 Oct 25 (pp. 205-219). Springer, Cham.
- [42] Kunnuji M, Eshiet I, Ahinkorah BO, Omogbemi T, Yaya S. Background predictors of time to death in infancy: evidence from a survival analysis of the 2018 Nigeria DHS data. *BMC public health*. 2022 Dec; 22 (1): 1-8.
- [43] Mekebo GG, Aga G, Gondol KB, Regesa BH, Woldeyohannes B, Wolde TS, Tadesse G, Galdassa A, Adebe KL, Ketema H. Why Babies die in the first 7 days after birth in Somalia Region of Ethiopia?. *Annals of Medicine and Surgery*. 2023 May 1; 85 (5): 1821-5.
- [44] Lamichhane R, Zhao Y, Paudel S, Adewuyi EO. Factors associated with infant mortality in Nepal: a comparative analysis of Nepal demographic and health surveys (NDHS) 2006 and 2011. *BMC public health*. 2017 Dec; 17 (1): 1-8.
- [45] Irana T, Mekebo GG, Diriba G, Sisay AL, Woldeyohannes B, Yohannes Z. Determinants of infant mortality in Oromia region, Ethiopia. *Annals of Medicine and Surgery*. 2023 Jun; 85 (6): 2791.